

All snake bites should be treated as potentially lethal

WARNING

- **DO NOT** try to catch or kill the snake.
- **DO NOT** allow casualty to move if possible. Bring help to the casualty.
- **DO NOT** wash or wipe venom off the skin. Hospitals are able to identify snakes from venom samples taken from the bite site.
- DO NOT cut bite site or area.
- **DO NOT** try to suck venom out of the bitten area.
- **DO NOT** use a constrictive bandage (i.e. arterial tourniquet).
- **DO NOT** remove bandage or splint once applied.



Code FFLI010

Manufactured by FASTAID 110 - 114 Old Bathurst Road enquiries please call Emu Plains, NSW 2750

www.fastaid.com.au

For consumer 1800 13 12 11 or +61 2 4728 8068



SIGNS & SYMPTOMS

After a person has been bitten, signs are not always visible and symptoms may only start to appear an hour or more later.



SNAKE BITE

- Puncture marks or scratches.
- ▶ Headache
- Nausea, vomiting, diarrhoea
- Double or blurred vision
- Breathing difficulties
- Drowsiness, giddiness, faintness drooping eyelids, problems speaking or swallowing
- Pain or tightness in chest or abdomen

SPIDER BITE

- ► Pain (moderate or severe)
- ► Nausea, vomiting and diarrhoea
- Excessive sweating

In addition to 'Spider Bite' Symptoms:

FUNNEL WEB SPIDER

- ► Abundant discharge of saliva
- Difficulty in breathing and muscular twitching
- ► Confusion leading to unconsciousness

RED-BACK SPIDER

- Severe local pain which increases and spreads
- ► Small hairs stand upright

OTHER SPIDER


- ▶ Inflammation or swelling
- Burning feeling
- Blisterina

WHAT SHO

SNAKE, FUNNEL-WEB

- **1. CALM THE CASUALTY.** Lie the casualty down. This will slow down the absorption of venom and reduce the effect of shock.
- 2. CHECK BREATHING AND CIRCULATION.

 If casualty is unconscious, follow DRSABCD
- **3. APPLY PRESSURE IMMOBILISATION BANDAGE.** Such as a crepe or conforming roller bandage around the bite site firmly.



Then apply a firm crepe or roller bandage over the whole limb starting at the toes or finger tips. Bandage needs to be firm as for sprain but not too tight.



4. IMMOBILISE THE LIMB USING A SPLINT.

If bitten on the leg, use the other as a splint. Use bandage to tie legs together or to hold splint in place. Keep the bitten limb level with the rest of the body to slow down absorption of the venom in the body.

ULD I DO?

MOUSE SPIDER BITES



- 5. CALL 000 or MOBILE 112 FOR MEDICAL HELP.
- **6. MONITOR CASUALTY.** Check at fingers or toes for circulation. If they lose consciousness or breathing fails **DRSABCD**

RED-BACK SPIDER

- APPLY WRAPPED ICE PACK to bitten area to reduce pain.
- **2. RAISE LIMB** to limit swelling.



3. SEEK MEDICAL HELP.

OTHER SPIDERS

- 1. WASH with water and soap.
- 2. APPLY WRAPPED ICE PACK to relieve pain or discomfort.
- 3. SEEK MEDICAL ADVICE.

BASIC LIFE SUPPORT FLOW CHART

CHECK FOR DANGER Hazards / Risks / Safety?



RESPONSIVE?
If not, call 000 for medical help

_

SEND FOR HELP
Call 000 (Triple Zero)
for an ambulance

OPEN AIRWAY.

Δ

Look for signs of life.

Signs of life = Conscious or responsive or breathing normally or moving.

GIVE 2 INITIAL BREATHS

If not breathing normally or properly.

GIVE 30 CHEST COMPRESSIONS

(Almost 2 compressions per second) then give 2 breaths.

ATTACH AED

(Automated External Defibrillator) as soon as available & follow its prompts.

•

CONTINUE CPR
until qualified personnel arrive
or signs of life return.

SNAKE & SPIDER IDENTIFICATION



WESTERN BROWN SNAKE (OR GWARDER)

Colour: from light brown to dark shade **Markings:** sometimes with crossbands

Nature: Highly venomous and extremely fast.

Location: WA, SA, NT, QLD, some of VIC



EASTERN BROWN SNAKE

Length: average 1.5-1.8m

Colour: from light brown to dark shade

Markings: sometimes with crossbands

Nature: timid but aggressive if provoked, extremely fast, highly venomous

Location:QLD, NSW, VIC, SA, arid parts of NT, WA (Kimberley region)



RED-BELLIED

Length: average 1.8
Colour: gleaming b

Markings: red lower

Nature: not very ag Location: East coa

river flats.



DUGITE SNAKE

Length: up to 2m

Colour: grey, green or brown

Markings: sometimes with crossbands

Nature: generally avoid humans, most active October-November,

highly venomous

Location: WA, coastal parts of west SA



DEATH ADDER

Length: up to 90cm

Colour: shades of grey through brown

Markings: irregular cross-banding

Nature: appears sluggish,

strikes with speed

Location: Throughout Aus, except

central deserts in SE NSW.



TAIPAN

Length: up to 3m

Colour: pale to dar

Markings: yellowis

Nature: extremely a

unprovoked attacks

Location: Northern



5-2m lack body r scales

BLACK

gressive st, swamps,



TIGER SNAKE

Length: up to 2.1m

Colour: pale grey to dark brown

Markings: yellowish cross-banding Nature: not especially aggressive,

but numerous and deadly

Location: Southern WA. SA. TAS. VIC.

NSW. Mainly Coastal Areas.



FUNNEL-WEB SPIDER

Length: 10 - 50mm

Colour: dark, ranging from

black to brown

Nature: extremely aggressive,

rear back to bite

Location: Eastern Coast of Australia



k brown n areas around head aggressive, given to , deadly venom

Australia



COPPER HEAD

Length: up to 180cm

Colour: varies from coppery mid-brown to yellowish, redish, grey and black

Nature: shy, retiring & prefer to escape rather than fight

Location: Temperate territories of Southern & Eastern Aus, well adapted

to cooler climates near water.



RED-BACK SPIDER

Length: 5-15mm

Colour: black body with prominent

red stripe on its abdomen

Location: throughout Australia